

Cobrexilin[®] 10

VETERINARY USE

Mineral supplement for the prevention and treatment of hypocuprosis in cattle.
Sterile suspension for injection.

Formula

Copper glycinate 10.0 g
Excipients q.s. 100.0 mL

Indications

It is indicated for the prevention and treatment of copper deficiency, primary or associated to molibdenosis in cattle.

Dosage and Administration

Administer by the subcutaneous route only in the lower part of the neck, near the union with the thorax.

Due to its low viscosity, 0.9 mm needles or similar can be used.

Shake well before using to ensure full resuspension of particles at the bottom of the bottle. The volume in the syringe should be accurately measured according to dose to be used.

Cobrexilin 10 may be combined with other treatments such as internal and external antiparasitics and others, but under no circumstance should the product be mixed with other products or medication in the same syringe.

Animals under 250 kg of bodyweight: 2 mL per animal. Repeat every 3 months.

Animals over 250 kg of bodyweight: 4 mL per animal. Repeat every 6 months.

First service heifers: 4 mL per animal, 1-2 months before service.

Open cows and bulls: 4 mL per animal, 1-2 months before service.

Pregnant cows: 4 mL in the last quarter of the pregnancy.

Important: Do not increase the recommended dose and do not shorten the intervals indicated in the treatment.

Package

Vial containing 250 mL.