

Maxillia[®]500 Soluble Powder

VETERINARY USE

Wide-spectrum antibiotic. Prescribed to combat infections caused by conditions caused by amoxicillin-sensitive infectious agents in the respiratory, genital-urinary, digestive systems and soft tissues and bacterial complications of viral conditions. To administer in drinking water.

Formula

Contents every 100 g of product:

Amoxicillin (as trihydrate) 50 g*
Excipients q.s. 100 g

*Equivalent to 57.4 g trihydrate amoxicillin.

Indications

Prescribed to treat: Actinobacilosis, Cystitis, Dermatitis, Tail Necrosis, Nephritis, Omphalitis, MMA Syndrome, Sinovitis by *Staphylococcus spp*, *Hæmophilus parasuis*, *Pasteurella multocida*, *Streptococcus pneumoniae*, *Streptococcus suis*, *Arthritis*, *Colibacillosis*, *Infectious Coryza*, *Infectious Enteritis*, *Chlostridial Diseases*, *Pasteurellosis*, *Salmonellosis* and other diseases caused by amoxicillin-sensitive microorganisms.

Dose and administration

Administer by mouth, mixed in drinking water, by virtue of 40 mg a day of Maxillia per kilogram of live weight, equivalent to 20 mg amoxicillin. Duration of treatment is 3 to 7 days, at the discretion of the Veterinarian.

The dose is obtained by preparing drinking water according to the following recommendations, based on the animal category and the evolution of the disease under treatment and average water consumption.

Poultry and pigs: 400 grs. Maxillia 500 Soluble Powder in water, every 10,000 kg. of live weight, a day for 3 to 7 days.

- For poultry, 0.266 g Maxillia per liter of water per day.
- For porcines: 0.70 g Maxillia per liter of water per day

Tip: Prepare the premix with the total amount of Maxillia to be administered in just 10 liters of water. Once the premix has been diluted, add the total water to be administered during treatment.

Prepare medicated water to be consumed in the next 6 to 8 hours. Do not store remainders for subsequent use.

Presentation

Plastic container for 1 kg.

Three-foil bag for 5 kg.

BROUWER