

# Meltra<sup>®</sup> Horses

USO VETERINARIO  
Venta bajo receta

Internal antiparasite paste for horses.

## Formula

Fenbendazole .....	18.75 g
Closantel .....	18.75 g
Excipients q.s. ....	100.00 mL

## Indications

For the treatment and control mature and immature stages of the following parasites in horses: *Parascaris equorum*, *Habronema sp*, *Oxyuris equi*. Large migratory strongyles: *Strongylus vulgaris*, *Strongylus equinus*, *Strongylus edentatus*. Small non-migratory strongyles: *Triodontophorus sp*, *Trichonema sp* and juvenile forms of *Gasterophilus nasalis* and *Gasterophilus intestinalis* (bots).

Meltra is indicated for the prevention of verminous aneurysm caused by migratory larvae of *Strongylus vulgaris*.

It can be used on animals of all ages and sex.

It has no effect on the fertility of males or the pregnancy of mares, or the fetus at any stage of the gestation.

It has neither embriotoxic nor teratogenic effects in foals.

It does not contain organophosphates.

## Dosage and Administration

It is administered orally: insert the tip of the dosing syringe into the horse's mouth and push the plunger slowly.

Due to its semisolid consistency, Meltra Equinos adheres itself to the buccal mucosa, avoiding spillage of the product and ensuring the proper dose.

The content of a 20 mL syringe of Meltra Equinos is sufficient for the treatment of a horse of 500 kg. To prepare the dose, use the scale of the plunger which starts in 100 and ends in 500 kilograms. These numbers indicate the weight of the horse to be treated.

Repetition of treatment:

To prevent verminous aneurysm, repeat treatment in 60 days.

To prevent juvenile forms of *G. nasalis* and *G. intestinalis*, treat every 60 days during spring and summer.

## Package

Disposable syringe containing 20 mL.