

Ophthalmologic infection preventive and healing treatment for cattle and sheep. Sterile solution.

Formula

Every 100 ml contains:

Tilmicosin phosphate	2 g
Hydrolyzed casein.....	2 g
Excipients qs.....	100 ml

Indications

Midribyl is indicated in treatment of infectious keratoconjunctivitis and other eye infections in cattle and sheep caused by the following microorganisms susceptible to Tilmicosin:

Moraxella bovis
Pasteurella multocida
Mycoplasma bovoculi
Staphylococcus aureus
Mycoplasma dispar
Streptococcus agalactiae
Chlamydia spp
Histophilus somni (Haemophilus somnus)
Actinomyces pyogenes
Actinobacillus pleuropneumoniae
Erysipelothrix rhusiopathiae
Mannheimia haemolytica
Clostridium perfringens
Fusobacterium necrophorum

Dosage and Administration

Applied in the treatment of infectious keratoconjunctivitis in cattle and sheep by means of a spray or dropper until the surface of the eye is appropriately covered; generally this is obtained by actuating the spray device 1 to 3 times dispensing 0.3 ml each time. The product should be applied every 12 hours and treatment should be continued during 4 days until symptom relief, though it can be prolonged according to the criterion of the acting Veterinarian.

Package

Package containing plastic 200 ml bottle with dosing spray device.