

External and internal antiparasitarian for bovines. Sterile solution for injection.

Formula

Doramectin 1 g
Inactive ingredients q.s. 100,0 mL

Indicaciones

Gastrointestinal Parasites

Ostertagia ostertagi: adults, L3, L4 (including inhibited states of *Ostertagia spp*); *Ostertagia lyrata*: adults and L4; *Haemonchus placei*: adults L3 and L4; *Trichostrongylus axei*: adults and L4; *T. colubriformis*: adults and L4; *T. longispicularis*; *Cooperia oncophora*: adults and L4; *C. pectinata*: adults, L4; *C. puntacta*: adults L4; *C. surnabada* (syn. *C. mcmasteri*); *Oesophagostomum radiatum*: adults, L3 and L4; *Nematodirus helvetianus*: adults; *N. spathiger*: adults; *Strongyloides papillosus*: adults; *Bunostomum phlebotomum*: adults, L3 and L4; *Trichuris spp*; *Thelazia spp*.

Pulmonary Parasites

Dictyocaulus viviparus: adult, L4, and inhibited states.

Ura (Torsalo, Nuche, Berne)

Dermatobia hominis, *Hypoderma bovis*.

Lice

Linognathus vituli, *Haematopinus eurysternus*, *Solenopotes capillatus*.

Mite

Psoroptes ovis (sin. *P. comunis variety bovis*); *Sarcoptes scabiei var. bovis*. It helps to control *Chorioptes bovis*.

Myasis

Cochliomya hominivorax.

Indications and Method of use:

Administer only SUBCUTANEOUSLY, in single dose at neck table.

The dose is 0.200 mg doramectin/kg of live weight, equivalent to 1 mL finished product per each 50 kg of live weight.

weight (kg)	dose (mL)	weight (kg)	dose (mL)
up to 50	1	301 to 350	7
51 to 100	2	351 to 400	8
101 to 150	3	401 to 450	9
151 to 200	4	451 to 500	10
201 to 250	5	501 to 550	11
251 to 300	6	551 to 600	12

To facilitate product use:

- When using multi-dose syringes, place a second needle at the cap to allow for air inlet, avoiding the vacuum which hampers handling. This is not necessary in case of automatic syringes which are provided with an air inlet valve.
- Protect the product from low temperatures. .

Package

50 and 500 mL white bottle.