

# Rolenal<sup>®</sup> Injectable

Analgesic and hypnotic opioid for dogs and cats. Sterile solution for injection.

## Formula

Tramadol ..... 50 mg  
Excipients q.s. .... 1 mL

## Indications

Tramadol is indicated as anesthesia premedication in order to improve intraoperative analgesia and enable patients to undergo a comfortable recovery. Its duration of action (between 6 and 8 hours) provides a significant analgesic effect in the immediate postoperative period. It is also indicated for the treatment of post surgical pain, being an effective analgesic after mastectomy and abdominal, orthopedic and spine surgeries. In addition, it is used in the treatment of acute pain in various syndromes, such as osteoarthritis and spinal pain.

## Dosage and Administration

The recommended dose is 1 to 2 mg/kg regardless of the route of administration. Therefore, we recommend the application of 0.1 mL every 2.5 to 5 kg of bodyweight.

The product can be used for the treatment of acute pain, as anesthesia premedication, as a painkiller during surgery and during the immediate postoperative period or for prolonged treatment. When treatment is prolonged, although oral administration is preferable, in some animals it may be more convenient to apply injections.

For these cases, we suggest an initial dose of 2 mg/kg every 6 hours for 24 - 48 hours later reduced to 1 mg/kg or administered at longer intervals of 8 hours, as the case requires. The duration of a complete treatment session must be determined by the acting veterinarian but for post-surgical pain usually only 3 to 5 days are necessary.

## Package

Vial containing 20 mL.