

Controls scabies; kills lice and flies. For use in cattle, horses, mules and swine. Emulsifiable liquid.

Formula

Cypermethrin	5.6 g
Diazinon	56.0 g
Excipients q.s	100.0 mL

Indications

Controls scabies and lice in cattle, horses, mules and swine. Kills flies: horn flies in cattle, flies in general.

Dosage and Administration

Spray on. To prepare correctly, pour Tipertox on to water, stirring until a homogeneous emulsion is obtained. Premixes are not necessary. Do not treat tired, weak or sick animals, or during times of the year when it is excessively hot. If it rains during the first hours after application, this must be repeated.

Against scabies: 145 mL every 200 liters of water. Apply 2 treatments with Tipertox 10 to 12 days apart. It is necessary to wet the animals thoroughly, especially in those parts of the body where there are more parasites. If the animals have crusted lesions these must be "brushed" with a brush or scraper until the scabs break up.

Against lice: 145 mL every 400 liters of water. One treatment is sufficient. Repeat in case of re-infestation.

Horn fly: 145 mL every 200 liters of water. Wet the animal's body completely. Repeat in case of re-infestations.

Control of flies and other insect species: roofs, walls, aisles between cages of laying hens and breeding swine, manure cones, floors: 145 mL in 200 liters of water.

Spray the masonry every 10 to 20 days. Repeat in case of re-infestations.

Package

Container with 145 mL.

