

Ambramicina lactancia®

VETERINARY USE

Antibiotic combination for intramammary application during lactation.

Formula

Cloxacillin (sodium) 0.20 g
Ampicillin (sodium) 0.075 g
Oil vehicle sufficient quantity for one syringe 10 mL

Indications

Prescribed to control mastitis produced by the following germs:

Gram (+) bacteria: *Staphylococcus aureus* (even in penicillin-resistant strains) *Streptococcus spp* (even *enterococcus*); *Corynebacterium pyogenes*; *Clostridium perfrigens*; *Cl. pseudotuberculosis*; *Listeria monocytogenes*; *Bacillus anthracis*.

Gram (-) bacteria: *Haemophilus influenzae*; *Escherichia coli*; *Salmonella spp*; *Shigella spp*; *Proteus mirabilis*; *P. morganii*; *Brucella spp*; *Neisseria spp*; *Bordetella pertussis*.

Target Species

Cattle during lactation.

Dose, method and way of administration

Cattle: One injector per quarter, that is 10 ml Ambramicina Lactancia®, equivalent to 75 mg sodium ampicillin and 200 mg sodium cloxacillin per quarter) every 12 hours. The complete treatment includes 3 doses.

Presentation

Package with 24 and 120 10-ml syringes.

BROUWER