

Meltra® Endectocide

VETERINARY USE

External and internal antiparasite drug for cattle. Sterile solution for injection.

Formula

Ivermectin 1.0 g
Excipients q.s. 100.0 mL

Indications

Antiparasitic activity:

Gastrointestinal Parasites

Ostertagia ostertagi: adults, L3, L4 (including inhibited stages of *Ostertagia spp*), *Ostertagia lyrata*: adults and L4, *Haemonchus placei*: adults, L3 and L4, *Trichostrongylus axei*: adults and L4, *T. colubriformis*: adults and L4, *Cooperia oncophora*: adults and L4, *C. punctata*: adults, L3 and L4, *Oesophagostomum radiatum*: adults, L3 and L4, *Nematodirus helvetianus*: adults, *N. spathiger*: adults, *Strongyloides papillosus*: adults, *Bunostomum phlebotomum*: adults, L3 and L4, *Toxocara vitulorum*: adults. Lung Parasites
Dictyocaulus viviparus (adults, L4 and inhibited stages).

Scabies Mites

Psoroptes ovis (*bovis variety*), synonym: *Psoroptes comunis var. bovis*.

Lice

Linognathus vituli, *Haematopinus eurysternus*, *Solenopotes capillatus*.

Human Bot Fly

Dermatobia hominis, *Hypoderma spp.*

Myiasis

Helps to control *Cochliomya hominivorax*.

Dosage and Administration

Administer subcutaneously in the front or back of the shoulder blade with 15-20 mm long, 16-gauge needle. Administer 1 mL of Meltra Endectocida every 50 kg of liveweight or 200 mg of Ivermectin per kg of liveweight.

Package

Vial containing 50 mL, 500 mL and 1 liter.

BROUWER