

Sintomicetina[®]

Broad-spectrum antibiotic for dogs and cats

Formula

Each multi-dose tablet contains

Azithromycin 200 mg

Excipients q.s..... 400 mg

Indications

Antimicrobial – Broad-spectrum antibiotic for the treatment of infections in soft and hard tissues, skeletal system, osteomyelitis and septic arthritis; respiratory tract infections, with an effective penetration into the lungs, larynx, trachea and nasal sinus; skin and skin-related infections, glandular secreting tissue, mastitis, metritis, ear infections, oral infections such as gingivitis, periodontitis, pulpitis, abscesses, cellulitis, and any infectious process caused by azithromycin-sensitive bacteria.

Taken once a day Azithromycin is particularly active against the following bacteria:

Infections of the respiratory tract caused by germs of the genus Haemophilus, Moraxella, or Streptococcus Infectious tracheobronchitis (kennel cough) caused by Bordetella bronchiseptica.

Pneumonia and bronchopneumonia caused or complicated by microorganisms of the genus Bordetella, Chlamydia, Haemophilus, Streptococcus, or Mycoplasma.

Oral infections caused by Streptococcus sp., Staphylococcus aureus and other common bacteria.

High respiratory tract infections caused by Streptococcus pyogenes as an alternative to treatment with Penicillin.

Pyoderma and infections of the skin caused by Staphylococcus aureus, Streptococcus pyogenes, or Streptococcus agalactiae.

External or middle ear otitis caused by germs of the genus Haemophilus, Moraxella, or Streptococcus.

Feline pneumonitis caused by Chlamydia psitaci.

Dosage and Administration:

Oral administration.

The total dose is 30 mg/kg of bodyweight.

Regular treatment: the total 30 mg/kg dose is divided into three intakes of 10 mg/kg each, with 24 hours intervals. One 200mg tablet every 20kg of bodyweight/day, during three consecutive days.

5 day treatment: one tablet every 20 kg of bodyweight during the first day, followed by half a tablet from the second day to the fifth day.

Single dose treatment: a tablet is administered every 6,6 kg of bodyweight (Gastric tolerance should be taken into account. In case of vomiting within an hour of intake, it is advisable to recommence treatment with the usual plan of 10 mg/kg of bodyweight for three days).

Package:

Box containing 6 multi-dose tablets.