

Triclacine®

VETERINARY USE

Internal antiparasite drug, indicated for the treatment of fascioliasis in cattle.
Aqueous suspension ready for use.

Formula

Triclabendazole 10 g.
Excipients q.s. 100 mL

Indications

For the treatment, prevention and control of both acute and chronic fluke. The product is effective against all stages of flukes, including early and immature stages of the trematodes. It is the product of choice for the treatment of acute and subacute outbreaks and for strategic control of *Fasciola spp.* infestations. It makes it possible to control infestations by the trematodes: *Fasciola hepatica* and *Fasciola gigantica* in cattle.

Dosage and Administration

Administer orally at a dose of 6 mL every 50 kg of bodyweight in cattle. This ensures a dose of 12 mg/kg of triclabendazole, the internationally recommended dose for this active principle in cattle.

In the treatment of acute outbreaks, one dose is enough to stop the epidemic cycle. Whether repeat treatment is necessary at 5 or 6 weeks should be decided by the acting veterinarian.

Strategic treatment programs should be designed specifically by the acting veterinarian on each facility. It is generally recommended to treat in late winter or early spring, so as to eliminate the adult *Fasciola spp.* (installed in the fall) and, thus, avoid vector infestation (*Lymnaea viatrix*; *L. truncatula*; *L. columella*). If the animals remain in the same facility, it is advisable to repeat the treatment in early summer. This is not necessary when animals are moved to winter pastures. In all cases, it is advisable to repeat the treatment in late summer – early autumn; later treatment may be necessary to eliminate parasites ingested in the spring and summer. These treatments must be supported by good management procedures such as field drainage, cleaning of canals, etc.

Use Restrictions: Pre-slaughter: 28 days.

Do not use in cows producing milk for human consumption.

Package

Backpack containing 2.5 liters.