

ACEDERM

FOR VETERINARY USE
Prescription medication

For use in veterinary medicine. Prescription medication. Hydrocortisone aceponate solution for topical administration. Antipruritic/anti-inflammatory agent. Dogs.

Acederm is formulated based on hydrocortisone aceponate: a corticoid of the diester family that is characterized by a rapid penetration through the stratum corneum of the skin to be metabolized in dermal structures. This kinetics in particular, its metabolism at the dermal level, allows to generate a powerful anti-inflammatory effect that minimizes local and systemic side effects.

Formula

Hydrocortisone Aceponate 0,584 mg
Excipients, q.s. 1 ml

Indications

Acederm is indicated for the treatment of inflammatory and pruritic dermatitis in dogs.

Route of administration, dosage and duration of treatment

The product is administered on the skin and does not require massages after its application. It must be sprayed at a distance of 10 cm from the area to be treated. The recommended dose per day is 1.52 Qg hydrocortisone aceponate per cm² of affected skin. This dose can be obtained by pressing the sprayer twice on the surface to be treated, equivalent to a square of 10 cm x 10 cm.

Daily administration for a period of 7 days is recommended. Depending on the origin of the dermatitis -inflammatory or pruritic- treatment may be continued at discretion of the Veterinarian.

If the animal is wet, both the skin and hair must be dry before treatment.

Avoid contact with eyes and mouth of the animal.

Contraindications

Do not use if there is a skin ulcer.

Precautions

In lieu of specific information, the risk-benefit ratio must be evaluated when using the drug in animals with Cushing's syndrome.

It is known that glucocorticosteroids stunt growth, so use in young animals (under 7 months) must be decided by the Veterinarian according to the risk-benefit ratio of the particular situation.

Use during pregnancy and lactation: safety of the veterinary medicinal product during pregnancy and lactation has not been demonstrated.

Use only in accordance with the benefit-risk assessment carried out by the Veterinarian.

In those cases in which the affected area involves a large part of the animal's body surface, the product must be used at the discretion and judgment of the Veterinarian; in these cases, periodic clinical studies are recommended.

Since the formulation may have a bitter taste, hypersalivation may occur if the animal licks immediately after application. This will disappear within some minutes, without the need to undergo any treatment.

Side effects

Transient local reactions may rarely occur at the application site (erythema).

Warnings

- Do not drink, smoke or eat during application.
- Do not spray on flame.
- Wash hands after application.
- Avoid contact with eyes, lips and mucous membranes. In case of contact with the eyes, wash with plenty of water.
- Do not store with food.

- Keep out of reach of children and pets.
- Avoid its ingestion. In case of accidental ingestion, call the Physician.
- Store in its original tightly-sealed container in a dry place, protected from direct sunlight and between 15°C and 30°C.

Dosage Form

Bottle with cap and spraying pump enclosed. It contains 20 ml that yield 153 applications.

 **senasa** - Certificate No. 17-107.

Manuf. Plant No. 4384 Made in Argentina

BROUWER S.A.

Technical Direction: Veterinary Physician

Rodolfo A. M. Perotti, License No. 2705

Dr. Rafael Bielsa 232/8 (C1427AZD) Buenos Aires.

Tel.: (54 11) 4555-6663.

www.brouwer.com.ar